




Preliminary studies on the quantification of the resilience of coastal ecosystems under anthropogenic pressures





Eric GOBERVILLE

in collaboration with
**Nicolas DJEGHRI, Loïc GUILLOUX, Philippe PONDAVEN,
Peggy RIMMELIN-MAURY and Paul TREGUER**

Master 2 internships
Lorène POURIAS (2024) and Lisa MOUQUE (2025)





Coastal marine ecosystems are more and more highly impacted





How European ecosystems have changed over the past decades?

Long-term changes in zooplankton and phytoplankton communities in European waters




Kendall statistic
4
0
-4





Attenuation
Climate index
Current
Mixed layer depth
Nutrients
pH
Precipitation
Salinity
Temperature
Wind




How French ecosystems have changed over the past decades?



Long-term changes in zooplankton and phytoplankton communities in European waters






Coastal marine ecosystems are more and more highly impacted



Future phytoplankton diversity in a changing climate


Stephanie A. Henson¹ , B. B. Cael¹, Stephanie R. Allen^{1,2,5} & Stephanie Dutkiewicz^{1,3,4}

2080-2100
(RCP8.5)
vs 2000-2020




Trajectories of the Earth System in the Anthropocene: The Hothouse Earth Pathway

Illustration of possible future pathways of the climate against the background of the typical glacial–interglacial cycles







Trajectories of the Earth System in the Anthropocene: The Hothouse Earth Pathway

Illustration of possible future pathways of the climate against the background of the typical glacial–interglacial cycles



Ordination trajectories and the ball-and-cup analogy



Ecosystem recovery and resilience

... the speed with which the system returns to its initial state following a disturbance (Holling)

Aim

Assess stability and resilience of coastal ecosystems by comparing present biodiversity with pre-industrial baselines

Method

DNA analyses from sediment cores (Bay of Brest, ~1,400 years) focusing on planktonic protists

Key findings


- WWII heavy-metal pollution → major community shifts
- 1980s–1990s agricultural pollution → further shifts in protist assemblages

Conclusion

No return to the Medieval baseline → irreversible shifts caused by cumulative anthropogenic pressures (war + agriculture)


Report

Sediment archives reveal irreversible shifts in plankton communities after World War II and agricultural pollution




How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

strong
low
mixing intensity





marine-dominated
open coastal
eutrophic



eutrophication




How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?







What factors control the temporal changes in oxygen concentrations in these coastal ecosystems?
Can these changes be modelled — and therefore predicted?



How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?







What factors control the temporal changes in oxygen concentrations in these coastal ecosystems? Can these changes be modelled — and therefore predicted?




How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

What factors control the temporal changes in oxygen concentrations in these coastal ecosystems?
Can these changes be modelled — and therefore predicted?






How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

How do local environmental conditions influence the spatio-temporal dynamics of phytoplankton groups?
Can we observe changes in recent years?





How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

How do local environmental conditions influence the spatio-temporal dynamics of phytoplankton groups?
Can we observe changes in recent years?




How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

How do local environmental conditions influence the spatio-temporal dynamics of phytoplankton groups?
Can we observe changes in recent years?





⇒ significant predictors
at different time scales



How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

How do local environmental conditions influence the spatio-temporal dynamics of phytoplankton groups?
Can we observe changes in recent years?



How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

How do local environmental conditions influence the spatio-temporal dynamics of phytoplankton blooms?
Can we observe changes in recent years?


Main findings

- Brest \Rightarrow Spring profile
- Roscoff \Rightarrow Clearer seasonal gradient
- Vilaine \Rightarrow Strong riverine influence

Conclusion

Confirms the hypothesis that ecological responses are locally structured

Spatial patterns



How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

How do local environmental conditions influence the spatio-temporal dynamics of phytoplankton blooms?
Can we observe changes in recent years?


Main findings

- Bacillariophyceae ⇒ Nutrient-rich spring conditions
- Dinophyceae, Euglenophyceae ⇒ Summer conditions
- Cryptophyta ⇒ Dispersal and ecological plasticity?

Conclusion

Helps identify the dominant ecological niches of each group

Phytoplankton patterns



How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

How do local environmental conditions influence the spatio-temporal dynamics of phytoplankton blooms?
Can we observe changes in recent years?


At seasonal scale

Components close to high phosphate concentrations
→ typical spring blooms


At long-term scale

→ Potential signals of climate change


Time-scale patterns



How can the resilience of coastal ecosystems under anthropogenic pressures be quantified?

* SORTING FEATURE COMING SOON
* CONTINUE TO AUCTION 

Contribution of plankton time series to quantifying the resilience of coastal ecosystems: study of the dynamics of trophic coupling between phytoplankton and zooplankton



Ehrlich & Gaedke (2020)



Thanks!!
Questions??

